

84641

S/020/60/133/005/020/034XX
C111/C222

Adams' Spectral Sequence

$K(z_{p,n})$.

3) Natural projections $p_j: \bar{X} \rightarrow X_j$ induce the epimorphisms $(p_j)_*$ for all $j \geq 0$.

Let $\alpha = \{X_i, \mathcal{P}_i, F_i, p_i\}$ and $\beta = \{X'_i, \mathcal{P}'_i, F'_i, p'_i\}$ be two p-systems. Let

$\alpha > \beta$ if for the identification of the groups $\pi_*(X), \pi_*(\bar{X})$ to $\pi_*(\bar{X}')$ it holds: $\ker(p_i)_* = \ker(p'_i)_*$, $i \leq n$ and $\ker(p_n)_* \subset \ker(p'_n)_*$.

Principal theorem 2: In the class of p-systems of a space X there exists one and only one system (up to the singular homotopic type) being maximal with respect to the introduced order.

Definition 3: The natural filtration of the p-space is the sequence of subspaces of \bar{X} :

$$\bar{X} = Y_0 \supset Y_1 \supset \dots \supset Y_3 \supset \dots,$$

where $Y_1 = p_1^{-1}(F_1)$, $\{X_i, \mathcal{P}_i, F_i, p_i\}$ is the maximal p-system of X and \bar{X} is the limit space.

Definition 4: The filtration of the homotopic and homological groups of the p-space X is called natural if it is generated by a natural filtration of X .

Card 3/4

84641

S/020/60/133/005/020/034XX
C111/C222

Adams' Spectral Sequence

Lemma 3: The elements $\alpha \in \pi_{m+n}(S^n X)$ and $E\alpha \in \pi_{m+n+1}(S^n X)$, where E - covering, have the same natural filtration if $m < n-1$.

Theorem 3: The filtration of the group $\pi_n^S(X)$ which is determined by the Adams sequence is identical with the natural filtration of this group.

The assertion of Adams on $E_\infty(S^0)$ (S^0 - zero-dimensional sphere) follows from theorem 3.

There is 1 non-Soviet reference.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet im. M.V.Lomonosova
(Moscow State University im. M.V.Lomonosov)

PRESENTED: May 13, 1960, by P.S.Aleksandrov, Academician.

SUBMITTED: May 11, 1960

Card 4/4

L 09189-67
ACC NR: AP7002800 SOURCE CODE: UR/0144/66/000/009/0995/1000

AUTHOR: Vinogradov, A. L.; Rizun, V. I.; Chebotarev, Ya. P.

ORG: none

TITLE: Automatic control system for asynchronous motors for a lifting apparatus for the stopping period, using controlled valves

SOURCE: IVUZ. Elektromekhanika, no. 9, 1966, 995-1000

TOPIC TAGS: thyristor, electronic rectifier

ABSTRACT: In order to provide a source of controlled voltage for dynamic braking in asynchronous drive apparatus, the authors have created small controllable rectifiers, operating on the phase control principle, using silicon thyristors. The requirements for accuracy and range of control are satisfied by using a phase controller, in which the heating signal is applied in the form of a pulse with a steep leading edge, which provides accurate control of the anode current. The main element of the phase controller is a shift register made of pulse logical elements. The supply pulses for the logic elements are formed by a pulse generator whose generation frequency is inversely proportional to the difference signal created by the difference between the assigned and actual speeds of the machine. Movement equations for the lift machine are presented. Orig. art. has: 3 figures and 20 formulas. [JPRS: 39,183]

SUB CODE: 13, 09 / SUBM DATE: 03Sep65 / ORIG REF: 006

Card 1/1 ⁴¹⁰

UDC: 621.313.333+621.375

8925-1688

VINOGRADOV, A.M.

Adam's spectral sequence. Dokl.AN SSSR 133 no.5:999-1002 Ag
'60. (MIRA 13:8)

1. Moskovskiy gosudarstvennyy universitet im. M.V.Lomonosova.
Predstavleno akad. P.S.Aleksandrovym.
(Sequences (Mathematics))

VINOGRADOV, A. M.

VINOGRADOV, A. M., Tekhnika i NAUMOVA, N. A. Arkh., OVCHINNIKOV, A. A. Arkh.

Nauchno-issledovatel'skiy institut arkhitektury zhillishcha Akademii Arkhitektury SSSR

Karkasno-shchitovyye i shchitovyye derevyannyye odnoetazhnyye doma zavodskogo
izgotovleniya massovogo tipa. Page 71

SO: Collections of Annotations of Scientific Research Work on Construction, completed
in 1950.
Moscow, 1951

VINOGRADOV, A.M.

Grabbing device for hoisting reinforced concrete rings.
Rats. i izobr.predl. v stroi. no.89:3-5 '54.(MLRA 9:6)
(Hoisting machinery)

VINOGRADOV, A.M.

Some homotopic properties of a space imbedding a circle inside
a sphere or ball. Dokl. AN SSSR 156 no. 5:999-1002 Je '64.
(MIRA 17:6)

1. Moskovskiy institut radioelektroniki i gornoy elektromekhaniki.
Predstavleno akademikom P.S.Aleksandrovym.

EOI'SHAKOV, V.M.; VINOGRADOV, A.M.; DOROKHOV, A.N.; PAZAROV, I.V.; MERTUNYAN,
A.K.; ROMANOV, A.A.; SEMEMOVSKIY, V.D.

Floors made of large rolled gypsum cement concrete panels. Stroi.
mat. 7 no.9:26-28 S '61. (MIRA 14:11)
(Floors, Concrete)

VINOGRADOV, A.N., kand.ekonomicheskikh nauk; LECHKIN, V.V., inzh.

Centralization and mechanization of inventory accounting for available freight cars. Vest.TSNII MPS 19
no.5:49-51 '60. (MIRA 13:8)
(Railroads--Accounts, Bookkeeping, etc.)
(Railroads--Freight cars)

VINOURADOV, A. R.

Methods for lowering the costs of house construction. Moskva, Gos. izd-vo arkhitektury i gradostroit., 1950. 65 p. (51-25004)

TH435.V5

VINOGRADOV, Andrey Nikolayevich; KAZANSKIY, B., red.; TELEGINA, T.,
tekhn.red.

[Planning the lowering of housing construction costs] Plani-
rovanie snizheniya sebestoimosti rabot v zhilishchnom stroitel'-
stve. Moskva, Gosfinizdat, 1959. 103 p. (MIRA 13:3)
(Construction industry--Costs)

VINOGRADOV, Andrey Nikolayevich; NAUMOV, I., otv.red.; BOROZDIN, B., red.izdva;
TELEGINA, T., ~~otv.red.~~

[Cost planning in construction; based on practice in housing
construction] Planirovaniye sebestoimosti v stroitel'stve; na
opyte zhilishhnogo stroitel'stva. Moskva, Gosfinizdat, 1962.
186 p. (MIRA 16:1)

(Construction industry—Costs)

VINOGRADOV, A.N.; PESKOVA, L.N., redaktor; VERINA, G.P., tekhnicheskii
redaktor

[Machine calculation in the railroad transport industry] Mekhani-
zatsiia ucheta na zheleznodorozhnom transporte. Moskva, Gos. transp.
zhel-dor. izd-vo, 1953. 295 p. [Microfilm] (MLRA 7:10)
(Railroads--Accounts, bookkeeping, etc.)
(Calculating machines)

Contact for graphite electrodes. A. N. Vinogradov. Russ. 32,485, Oct. 31, 1933. The contact of low-melting metal, cast as a bead that surrounds the end of the graphite electrode, is placed in a jacket of a high-melting metal to prevent the low-melting metal from running off the electrode.

40. ASM-56A METALLURGICAL LITERATURE CLASSIFICATION

Apparatus for electrodepositon of magnesium. A. N. Vinogradov. Russ. 22,727, Oct. 31, 1963. For electrolysis of fused $MgCl_2$ the anodes and cathodes are superimposed in a number of horizontal rows forming vertical groups of electrodes of different polarities. Solid nonconducting rods are placed at the joints of these rows for the separation of the electrode products and for the withdrawal of the separating Mg under rods placed above the cathodes.

AND S. A. DETAILING LITERATURE CLASSIFICATION

Apparatus for electrodeposition of magnesium. A. N. Vinogradov and A. M. Krupnikov. Russ. 33,000, Dec. 31, 1953. The app. for electrolysis of fused chloride has horizontal anodes placed below the cathodes so arranged that the distance between the electrodes can be regulated and the consumption of the anodes compensated. The anodes are of triangular or trapezoidal cross section, and the cathodes are arranged parallel to the spaces left between the anodes.

ASR SLA METALLURGICAL LITERATURE CLASSIFICATION

VINOGRADOV, A.N.; LIVSHIN, G.L.; OBRAZTSOVA, R.I.; TULUPOV, L.P.;
Prinimali uchastiye: RAZORENOVA, L.K., inzh.; DUBINKINA,
L.I., inzh.; PODGORYKH, A.L., inzh.; LAVRENT'YEV, M.V.,
retsenzent; MINAKOV, A.D., retsenzent; NESTEROV, Ye.P.,
retsenzent; STEFANOV, N.Ya., retsenzent; USHAKOV, P.S.,
retsenzent; KRISHTAL', L.I., red.; KHITROVA, N.A., tekhn.
red.

[Calculating machines in accounting, planning and administration in railroad transportation] Vychislitel'naya tekhnika v uchete, planirovanii i upravlenii na zheleznodorozhnom transporte. [By] A.N.Vinogradov i dr. Moskva, Transzheldorizdat, 1963. 407 p. (MIRA 17:2)

SKLYUYEVA, M.A.; VINOGRADOV, A.N.; FEDIUNINA, Ye.N. (Moskva)

Education and postgraduate training of medical personnel. Sov.
zdrav. 21 no.1:85-96 '62. (MIRA 15:2)

1. Iz Otdela meditsinskoy statistiki (nachal'nik G.F.Tserkevnyy)
Ministerstva zdravookhraneniya SSSR.
(MEDICINE STUDY AND TEACHING)
(MEDICAL STATISTICS)

VINOGRADOV, Aleksandr Nikolayevich; CHERNOMORDIK, D.I., prof., retsenzent;
PETROV, A.I., red.; USENKO, L.A., tekhn. red.

[Why it is necessary to have separate accounting in every railroad department] Zachem nuzhen uchet v kazhdom zheleznodorozhnom predpriyatii. Moskva, Vses. izdatel sko-poligraf. ob"edinenie M-va putei soobshcheniia, 1961. 49 p. (MIRA 14:10)
(Railroads—Accounts, bookkeeping, etc.)

ALEKSEYEV, V.N.; VINOGRADOV, A.N.; kand.ekon.nauk; VLADIMIROV, V.A.; inzh.;
KOCHETOV, I.V., prof.; doktor ekon.nauk; MINAKOV, P.P.; POTAPOV,
I.A.; ROMANOV, M.P., dotsent, kand.ekon.nauk; SPENGLER, Ye.N.,
kand.ekon.nauk; SHITOV, A.V.; SHUKHATOVICH, I.M.; YAKUBOV, L.S.;
IVLIYEV, I.V., red.; KRISHTAL', L.I., red.; KOCHETOV, I.V., prof.,
doktor ekon.nauk, nauchnyy red.; IVANOV, A.P., nauchnyy red.;
BOBROVA, Ye.N., tekhn.red.

[Statistics and bookkeeping in railroad transportation; manual]
Statistika i bukhgalterskii uchet na zheleznodorozhnom transporte;
spravochnik. Moskva, Vses.izdatel'sko-poligr.ob"edinenie M-va
putei soobshcheniia, 1960. 485 p. (MIRA 14:3)
(Railroads--Accounts, bookkeeping, etc.)
(Railroads--Statistics)

ca

11 c
DEAN JBR.

Occurrence of manganese in insects. II. A. P. VINIKRAJIV AND M. V. NER-
TSEVA. *Compt. rend. acad. sci. U. R. S. S. 1910A*, No. 6, 127-32; cf. *Compt. rend.
acad. Sci. U. R. S. S. 1929A*, No. 10, 227.—Mn was detd. in a large no. of insect species.
The av. % of Mn based on live wt. was 10^{-4} , the extreme values being 2×10^{-4} and
 8.3×10^{-4} . *Formica rufa* has the max. content. The amt. of Mn depends somewhat
on the character of the food, is independent of sex and varies inversely with the am-
t. of gaseous exchange in the insect, those always in motion having the smallest Mn con-
tent. With regard to distribution in the body, in *Formica rufa* the head contains an
av. % of 2.8×10^{-4} , middle portion 9×10^{-4} , posterior portion 3×10^{-4} based on the
dry wt. The latter part of the insect contains the major portion of the alimentary tract
and also the gland furnishing the poisonous secretion. Its high Mn content is not
due to food residues in the tract since the food of these insects contains less Mn and
more Fe than the whole insect. As Mn plays a catalytic role in some fermentations,
and as HCO_2H probably results from the action of an oxidative enzyme upon carbohy-
drate, the high Mn of the posterior part may be related to HCO_2H production.
LAWIN W. BERRY

ASH-SLA METALLURGICAL LITERATURE CLASSIFICATION

112

Ca

Occurrence of vanadium in marine organisms. A. P. VINOGRADOV. *Compt. rend. acad. sci. U. R. S. S.* 1930A, No. 17, 465-7. -- *Phallusia obliqua* and *Ciona intestinalis* contain V; *Molgania affinis* and *Cucumaria frondosa* do not. It would seem that all Ascididae have a V contg. blood pigment. *Phallusia obliqua* contains 0.0002% of the live wt. The quant. method follows: Fuse the ash with Na_2CO_3 and KNO_3 ppt. its vanadate from the aq. soln. of the melt, dissolve the ppt. and titrate the V, reduced by 80% with K_2SnCl_6 .

LUDWIG W. HURT

ASH, SLA METALLURGICAL LITERATURE CLASSIFICATION


CA 11/D

PROCESSES AND PROPERTIES INDEX

The elementary chemical composition of some species of the genus *Lenno* as a species characteristic. V. I. VORNADSEKH AND A. P. VINOGRADOV... *Compt. rend. acad. Sci. U. R. S. S. 1631A*, No. 6, 148-52 -- A preliminary paper presenting the results of quant. analyses of *Lenno polyrhiza*, *L. minor*, *L. trisulca* and *L. gibbs* for H, O, C, N, Ca, K, Cl, Mg, P, Na, S, Si, Mn and Fe. Al, Ti, V, Ra, Sr, Ba, Rb and Cu are also present, but their detn. has not been completed. The results indicate that the elementary compn. can be considered a species characteristic in the same sense as the morphological characters.

LEWIS W. BUTZ

ASB. S.L.A. METALLURGICAL LITERATURE CLASSIFICATION



11J

Ca

Trudy,
Dobro

Vanadium in certain organisms. A. P. VINOGRADOV. *Trudy, Inst. biokhimi.*
1958, No. 11, R. S. S. 2, 1-7 (1952); cf. C. A. 26, 2138. V was detected in 0.1 g. of ash
of *Ciona intestinalis* L. and *Ascidella aspersa* Müll. Ash from *Sarcobolylloides aurea*
(3 g.) gave a hardly detectable, but pos. reaction for V. Ash of *Pyura* and *Sipia* (10 g.)
gave no reaction for V. *Ciona intestinalis* contained $1.3 \times 10^{-4}\%$ of V in the
living tissue and $4 \times 10^{-4}\%$ in the ash. *Ascidella aspersa* contained $4.9 \times 10^{-4}\%$ in
the living tissue and 0.1% in the ash. Detn. of V in 13 species of *Aurida* made by vari-
ous workers are given. V was not detected in *Malpasia affinis* and *Cucumaria frondosa*.
B. S. IVANOFF

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

1958: 23417V

1958: 23417V

CA 11-5

PROCESSES AND PROPERTIES INDEX

The chemical composition of plankton. I. Analysis of plankton from the Katerininski pond at Detoké Selo. A. Vinogradov. *Trav. lab. biogéochim. acad. sci. U. R. S. S. I.*, 33:44; *Neues Jahrb. Mineral. Geol., Referate II*, 1950, (NM-1).—Nineteen organisms identified are listed. The mean H₂O content was 80.77%; ash content of the living matter 1.01%, of the dried material 11.28%. Fifteen chem. elements were detd. in the living and dried material. J. P. Schauer

ASS-114 METALLURGICAL LITERATURE CLASSIFICATION

150000 151 000 152

150000 151 000 152

150000 151 000 152

CA

8

PROCESSES AND PROPERTIES INDEX
EST AND JMC CODES

The origin of iodine and bromine in oil-bearing waters.
A. P. Vinogradov. *Compt. rend. acad. sci. U. R. S. S.*
[N. S.], 7, 214-15 (in English) 210(1934).—The oil-
bearing waters of the Caucasus and the marine muds of the
Black, Barents and Kara Seas contain 0.003-0.02% I and
0.0025-0.007% Br. The origin of the I is in the fossil
food of marine bacteria. P. H. Rathmann

DOK. AN SSSR

AISI-SLA METALLURGICAL LITERATURE CLASSIFICATION


STANDARD PA STANCOB MAP INC JOE BRILLSTONE
VIZOR GMAHVV VIKASIT SHG QWO SAS

1ST AND 2ND ORDERS																										PROCESSES AND PROPERTIES INDEX																									
COMMON ELEMENTS																										METALLURGICAL LITERATURE CLASSIFICATION																									
<p>Distribution of vanadium in organisms. A. P. Yung, <i>sydney. Compt. rend. acad. sci. U. R. S. S.</i>, 194-8 (in English 404-9)(1934).—The concn. of V in most terrestrial and marine animals and plants is about 0.13%. <i>Aricidia</i> (Tunicate) is unusually rich in V (2×10^{-3} to 1.5%). Marine muds are the probable source of V. Oils contg. V may be formed from <i>Aricidia</i>. W. F. B.</p>																																																			

VINOGRADOV, A. P.

"The Elemental Chemical Composition of Microorganisms and D. I. Mendeleyev's Periodic System of Elements," Works of the Biogeochemical Laboratory of the Academy of Sciences USSR, Vol. 3, 1935.

1ST AND 2ND ORDERS										PROCESSES AND PROPERTIES INDEX									
<p>Biogenic migration of rare elements. A. P. Vinogradov. <i>Trans. Far East. Sci. Acad. USSR</i> 1935, A, 64. There appears to be a tendency toward the accumulation of elements of uneven at. no. in soil compared with the compn. of the earth's crust. Elements which show greatest biogenic migration are those which most easily form sol. compds. B. C. A.</p>										<p>8</p>									
<p>ASD-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>										<p>12</p>									



PROCESSING AND REPORTING																									
1ST AND 2ND EDITIONS													3RD AND 4TH EDITIONS												
1ST AND 2ND EDITIONS													3RD AND 4TH EDITIONS												
<p><i>11E</i></p> <p>The elementary chemical composition of marine organisms. A. P. Vinogradov. <i>Trav. lab. biogeochem. acad. sci. U. R. S. S. R.</i> 63-278 (1935); cf. C. A. 28, 1728. A treatise on the occurrence and importance of chem. elements in various species of marine organisms and their surroundings. N. N. Menshik</p>																									
<p>ASAC-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																									

VINOGRADOV, A. P.

Vanadium in the petroleum and bitumens of the U.S.S.R. A. P. VINOGRADOV and G. G. Bergman (Compt. rend. Acad. Sci. U.R.S.S., 1935, 4, 349-352; cf. A., 1934, 1380). The distribution of V in the petroleum and bitumens of the U.S.S.R. has been determined. Deposits of the Devonian period contain the highest and those of the Jurassic period contain the lowest % of V. From much of the data it appears that V occurs as the % of org. matter in the rock, in Ishma asphaltite. The view that V passes into the petroleum from aq. solution in presence of H_2S , the V being held as V_2S_5 , is considered disproved. It is probable that the V comes from the mud of continental sea basins in which V has been conc. by marine organisms. B. S. B.

SUBJECT		PROCESSES AND PROPERTIES INDEX	
ca	<p>The origin of vanadium in petroleum and hard bitumens. A. P. Vinogradov. <i>Akad. V. I. Vernadskomu k 50-letiyu Nauch. Deyatelnosti</i> 1, 115-67 (1960); <i>Chem. Zvezdy</i>, 1960, 1, 1711; <i>U. S. A. J.</i>, 1965. The occurrence of V in the ash of petroleum and hard bitumens in various parts of the earth is discussed. The V content of the ash of crude petroleum and bitumen reaches 80-90% and varies widely for samples from the same deposit. If the V content is calculated for the product as a whole and not for the ash, then rather uniform values are obtained. The V content, therefore, is connected with the org. portion of the petroleum and bitumen, according to Treloar the V is combined in the form of V-porphyrin compounds. V-contg. petroleum are characterized by a considerable content in asphalt and S. As to the origin of the V in petroleum and bitumen, it is pointed out that the V is always associated with Ni and the view is expressed that it comes from the sludge of the continental seas, in which it was present since the beginning of the formation of the petroleum, and together with the Ni present it extensively influenced the conversion of the org. matter into petroleum. The high S content of V-contg. petroleum is explained as due to the favorable influence of the V on the processes of oxidation of org. matter by sulfates. On the basis of geological data, chem. analyses, etc., it is shown that the hard, V-contg. bitumens are to be regarded as derived of petroleum, from which they have been formed by polymerization.</p> <p>ST. G. Moore</p>	2	
<p>ASH-SCA METALLURGICAL LITERATURE CLASSIFICATION</p>			

The elementary chemical composition of marine organisms. H. A. V. Vinogradov. *Izv. Akad. Nauk SSSR, ser. Khim. Nauk*, 1959, No. 1, p. 225 (103); cf. *Ch. 28, 1728; 29, 1807*.—Nyne *rep.* reviews on the compn. of various species of Bryozoa, Brachiopoda, Vermes, Enteropneusta, Echinodermata, Mollusca, Arthropoda, Tunicata, and of hemoglobin, hemocyanin and other respiration pigments of invertebrates comp. metals. Most of the data are assembled in tables. Only authors' names and yr. of publication are given. T. Laanes

Trudy. 15
 Lib. Disposition.
 All done.

A 5 D. 3 L A METALLURGICAL LITERATURE CLASSIFICATION

1ST AND 2ND ORDERS										PROCESSES AND PROPERTIES INDEX										100 AND 4TH ORDERS									
<p>Fluorine content in the rivers of the Union in connection with the occurrence of mottled enamel disease). A. P. Vinogradov, V. V. Danilova and L. N. Shivanov. <i>Chem. Acad. Sci. U. R. S. S. 16, 361-4 (1957) (in English).</i> The F content of various portions of ten large rivers in the Soviet Union does not exceed 0.2 p. p. m. A higher F content was found in waters located in the regions of the Khibiny apatite deposits. H. J. Prebula</p>																													
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																													
FROM STUDY										SEARCHED										SERIALIZED									
1000000000										1000000000										1000000000									
1000000000										1000000000										1000000000									

115

DATE: 11/1/58

Ca

Manganese in insects (Formicidae) III. The chemical composition of organisms as a specific character. P. Vinogradov. *Compt. rend. acad. sci. U. R. S. S. 14*. 1957-9 (1957) (in English); cf. C. A. 25, 1562. The Mn content of ants belonging to the subfamily *Camponotinae* (stingless) is about 0.01%; the stinging subfamilies, *Myrmicinae* and *Ponerinae*, have approx. 0.001% Mn. The Mn content of ants coincides with their membership in definite subfamilies. The Mn content is different and characteristic for the various genera of *Formicidae*, whereas it is uniform for the species of *Acrididae*. The peculiar distribution of Mn within these families indicates that families do not have equal significance as taxonomical units.

II 1. Prehuda

ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION

10000 11000 12000 13000 14000 15000 16000 17000 18000 19000 20000 21000 22000 23000 24000 25000 26000 27000 28000 29000 30000 31000 32000 33000 34000 35000 36000 37000 38000 39000 40000 41000 42000 43000 44000 45000 46000 47000 48000 49000 50000 51000 52000 53000 54000 55000 56000 57000 58000 59000 60000 61000 62000 63000 64000 65000 66000 67000 68000 69000 70000 71000 72000 73000 74000 75000 76000 77000 78000 79000 80000 81000 82000 83000 84000 85000 86000 87000 88000 89000 90000 91000 92000 93000 94000 95000 96000 97000 98000 99000

1ST AND 2ND ORDER										PROCESSES AND PROPERTIES INDEX										3RD AND 4TH ORDER									
<p>LA</p> <p>Analytical chemistry of rare elements in small concentrations. I. P. Alimarin and A. P. Anisimov. <i>Zhurnal. V. 8, 888-891 (1934). A discussion.</i> C. H.</p>																													
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p>																													
100000 00										100000 000 000 000										100000 000 000 000									
00 00 00 00 00 00 00 00 00 00										00 00 00 00 00 00 00 00 00 00										00 00 00 00 00 00 00 00 00 00									

VINOGRADOV, A. P.

Biogeochemical provinces and endemics. A. P.
Vinogradov (Compt. rend. Acad. Sci. U.R.S.S., *1938, 18, 283-286*).--The influence of the chemical
characteristics of soils, notably of excess or
deficiency of biologically active constituents
on the flora and on disease in plants and animals
is discussed. A.G.P.

LIST AND 77. CATEGORIES																										PROCESSES AND PROPERTIES, ETC.																									
COMMON ELEMENTS																										METALLURGICAL LITERATURE CLASSIFICATION																									
<p>Geochemistry and biochemistry. A. P. Vinogradov. <i>Ispolki Khim.</i> 7, 645 (1958). A review chiefly of the Soviet literature. V. discusses the chem. compn. of organisms from the point of view of the wide variety of elements (if not all) present, and the possible relations of these quantities to org. evolution as related to the various ams. of different elements available in different geol. epochs. Many biochem. and geochem. data on the compn. of sea water, of various protozoa, diatoms, plankton, etc., are given. Other tables give the H, O, C, N, S, P, K, Ca, V, I, Li, Cl and F in fresh water and marine plants, in various mollusca, algae, ascidians, etc., as well as in petroleum and bitumens, silt and soils, rivers, and lakes of the U. S. S. R. E. H. Rathmann</p>																																																			

PERCEIVED AND PROPERTIES INDEX																									
COMMON ELEMENTS													COMMON ELEMENTS												
<p>Iodine in marine muds. Origin of iodine- and bromine-containing waters in petroliferous regions. A. P. Vinogradov. <i>Trav. lab. biogéochim. acad. sci. U. R. S. S. R.</i>, 10 32 (in English, 33-40) (1939).--The dried muds of the Caspian, Aral, Black, Barents, White, Kara and Okhotsk seas contain 0.001-0.005% I, the fine-grain mud fraction about 0.05% of this amt. At different levels of mud strata the amt. of I depends upon the content of org. matter. The data indicate a possible connection between the concn. of I (and Br) in the water over petroliferous beds and its concn. in the mud and mud water. T. Lianes</p> <p><i>Trudy Biogeochem. Lab. 1939 558p.</i></p>																									
ASB-SEA METALLURGICAL LITERATURE CLASSIFICATION																									
COMMON ELEMENTS													COMMON ELEMENTS												

CA

Investigations of the biogeochemical regions in connection with their economic importance. A. P. Vinogradov. *Vestnik Akad. Nauk S. S. S. R.* 1939, No. 10, 109-17. *Khim. Referat. Zhur.* 1940, No. 7, 10. — The av. content of all chem. elements in each geochem. system (soils, water, etc.) is const. The org. universe developed about each of these systems. Deviations from the ordinary av. composition of soils, water, etc. caused peculiar biol. reactions of the flora and fauna in these regions and produced new morphol. changes in the species and pathol. processes (endemic). Regions with insufficient or excessive amounts of certain elements or groups of elements are characterized by the development of endemic diseases affecting plants and animals (mainly the young organisms). There are known biogeochem. regions with insufficient contents of H, I, P, B, Se, P, Li, Be, Mn, Mg, Ca, Sr, Cu, Co, Fe, Zn, Ni, Mo, etc., in soils and water. Methods against such endemic diseases in regions consist in regulating the contents of the single microelements in the food of plants and animals (I for goiter, Co for anemia of sheep, B for beriberi diseases, Cu for diseases of grain plants, etc.).

W. K. Henn

ASB-51A METALLURGICAL LITERATURE CLASSIFICATION

VINOGRADOV, A. P.

Geochemical investigation of the spreading
of the Urov disease. A. P. VINOGRADOV (Compt.
rend, Acad. Sci., U.R.S.S., 1939, 23 64--67).
An investigation of the drinking-H₂O in regions
where the Urov disease (akin to rickets) is
prevalent has been carried out. "Affected"
waters (i.e., those in the region where the
disease is prevalent) contain much less mineral
matter, and particularly Ca, than "healthy"
waters. A.J.M.

Biogeochem. Rab. Mhr. Acad. Sci. c1939-

VINOGRADOV, A. P.

"The Conference on the Chemistry of Isotopes."

Zhur. Fiz. Khim., Vol. 14, No. 5-6, 1940.

112

21

The cause of the absence of calcareous (CaCO_3) skeletons in pre-Cambrian invertebrates. A. P. Vinogradov. *Compt. rend. acad. sci. U. R. S. S.* 27, 232-5 (1940) (in English).—The absence of calcareous skeletons in pre-Cambrian invertebrates was due to the sea water being unsatd. with CaCO_3 , owing to a higher CO_2 content both in air and water. The formation of calcareous skeletons in invertebrates occurred under conditions of satn. (or supersatn.) of sea water with CaCO_3 . At present the process of formation of these skeletons is controlled by satn. of water with CaCO_3 . The photosynthesizing pre-Cambrian algae, consuming CO_2 and bicarbonate, created conditions for satn. with CaCO_3 while they were depositing CaCO_3 in their tissues. Felix Saunders

Biogeochemical Lab; Acad. Sci., - C 1940-

ASAC-SLA METALLURGICAL LITERATURE CLASSIFICATION

SECTION	SUBSECTION	TERMINOLOGY	ANALYSIS	SYNTHESIS	OTHER
1	1	1	1	1	1
2	2	2	2	2	2
3	3	3	3	3	3
4	4	4	4	4	4
5	5	5	5	5	5
6	6	6	6	6	6
7	7	7	7	7	7
8	8	8	8	8	8
9	9	9	9	9	9
10	10	10	10	10	10
11	11	11	11	11	11
12	12	12	12	12	12
13	13	13	13	13	13
14	14	14	14	14	14
15	15	15	15	15	15
16	16	16	16	16	16
17	17	17	17	17	17
18	18	18	18	18	18
19	19	19	19	19	19
20	20	20	20	20	20
21	21	21	21	21	21
22	22	22	22	22	22
23	23	23	23	23	23
24	24	24	24	24	24
25	25	25	25	25	25
26	26	26	26	26	26
27	27	27	27	27	27
28	28	28	28	28	28
29	29	29	29	29	29
30	30	30	30	30	30
31	31	31	31	31	31
32	32	32	32	32	32
33	33	33	33	33	33
34	34	34	34	34	34
35	35	35	35	35	35
36	36	36	36	36	36
37	37	37	37	37	37
38	38	38	38	38	38
39	39	39	39	39	39
40	40	40	40	40	40
41	41	41	41	41	41
42	42	42	42	42	42
43	43	43	43	43	43
44	44	44	44	44	44
45	45	45	45	45	45
46	46	46	46	46	46
47	47	47	47	47	47
48	48	48	48	48	48
49	49	49	49	49	49
50	50	50	50	50	50
51	51	51	51	51	51
52	52	52	52	52	52
53	53	53	53	53	53
54	54	54	54	54	54
55	55	55	55	55	55
56	56	56	56	56	56
57	57	57	57	57	57
58	58	58	58	58	58
59	59	59	59	59	59
60	60	60	60	60	60
61	61	61	61	61	61
62	62	62	62	62	62
63	63	63	63	63	63
64	64	64	64	64	64
65	65	65	65	65	65
66	66	66	66	66	66
67	67	67	67	67	67
68	68	68	68	68	68
69	69	69	69	69	69
70	70	70	70	70	70
71	71	71	71	71	71
72	72	72	72	72	72
73	73	73	73	73	73
74	74	74	74	74	74
75	75	75	75	75	75
76	76	76	76	76	76
77	77	77	77	77	77
78	78	78	78	78	78
79	79	79	79	79	79
80	80	80	80	80	80
81	81	81	81	81	81
82	82	82	82	82	82
83	83	83	83	83	83
84	84	84	84	84	84
85	85	85	85	85	85
86	86	86	86	86	86
87	87	87	87	87	87
88	88	88	88	88	88
89	89	89	89	89	89
90	90	90	90	90	90
91	91	91	91	91	91
92	92	92	92	92	92
93	93	93	93	93	93
94	94	94	94	94	94
95	95	95	95	95	95
96	96	96	96	96	96
97	97	97	97	97	97
98	98	98	98	98	98
99	99	99	99	99	99
100	100	100	100	100	100

CA

Copper content of various soils (the origin of the so called reclamation disease of cereals). A. P. Vinogradov. (Compt. rend. acad. sci. R. S. S. 27, 1092 8/1940) (in German). Lack of Cu in newly cultivated peat, sand and marsh soils may be the cause of the disease of cereals. Chem. and spectroscopic detns. were made of the Cu content of marsh soils of White Russia and various other soil types of the Soviet Union. The soil samples were incinerated in the presence of H_2SO_4 . SO_3 was removed by treatment with HF . The ppt. was dissolved in dil. HCl , the insol. ppt. was melted and likewise dissolved. In the dil. NH_4 soln. Cu was pptd. by means of rubenic acid. The Cu ppt. was carefully heated in the presence of H_2SO_4 in quartz retorts and dissolved. In the soln. Cu was detd. by the method of Spacu. The soil samples the Cu content of which had thus been detd. were used as standards in the spectroscopic analysis of the other soil samples. The Cu content of such soils is low (usually 2×10^{-4} to 1×10^{-3}). Marsh soils fertilized with Cu showed only a small increase of the Cu content. Cu is not stored by the org. substance of marshes. The Cu content of the upper humus layers of the Ukrainian wheat belt is somewhat higher than that of the lower layers. The Russian "red earths" are high in Cu. A. H. Kravtch.

Biochemical Lab. Mbu, Acad. Sci. c1940.

ASH 55.4 METALLURGICAL LITERATURE CLASSIFICATION

Isotopic composition of waters in metamorphic rocks and minerals. V. I. Vernadsky, A. P. Vinogradov, and R. V. Tsia (Copt. rend Acad. Sci. U. R. S. S., 1941, 31, 573-576).--Chlorites, talcs, and serpentines of different deposits and geological ages were used. H₂O was distilled from them at 1145°, and its δ determined. In almost all cases the δ was > that of standard purified tap-H₂O. In two cases, the greater δ was shown to be due entirely to ¹⁸O, the D content being < that of standard H₂O. A. J. W.

Isotopic composition of oxygen of different origin. A. P. Vinogradov and R. V. Teis (*Compt. rend. Acad. Sci. U.R.S.S.*, 1941, 28, 490—493).—Determinations of the δ of H_2O synthesised from H_2 (produced by the action of HCl on Zn) and O_2 from various sources indicate that O from the atm. is heavier than that from H_2O whilst that evolved under the action of light by *Helodora canadensis* growing in tap- H_2O containing 0.1% of $NaHCO_3$ is intermediate. The O in CO_2 produced by the action of HCl on $NaHCO_3$ is heavier than atm. O . J. W. S.

Biogeochem. Ab., *Mus. Acad. Sci.* c1941-

Decomposition of kaolin by diatoms. A. P. VINOGRADOV
AND K. A. BOICHENKO. *Compt. rend. acad. sci. U.R.S.S.*,
37, 135-38 (1942) (in English); *Chem. Abstracts*, 37, 5673
(1943).—Nacrite crystals (Simferopol, Russia) of 20 to 50 μ
were attacked by the slime secreted by diatoms. This
slime is composed of a pectinous substance. In the late
stage of the process, the nacrite crystals appear to exfoliate,
alumina hydrate is released, and the shells of former
nacrite crystals remain. Bacteria mixed with the diatoms
have no direct effect on the nacrite.

Biogeochem. Lab.: Acad. Sci., 1942.

CA

Geochemistry of the scattered rare elements in sea water. A. P. Vinogradov, *Uspekhi Khimii* 13, 3-31 (1944). - Review. Numerous data on the distribution of I, P, As, B, Li, Rb, Sr, Fe, Mn, Cu, Ni, Co, other heavy metals, Mo, Ag, Au, Pt elements, Yt, Sc, rare earths, and Ra and related elements in various oceans at different depths and times are shown in 20 tables. Great variations occur depending on the origin of the ocean waters and on the depletion of various elements by biol. and geochem. causes. The I and B cycles and the U \rightarrow Ra relations in nature are shown. 197 references. F. H. Rathmann

Ca

POSSIBLES AND PREPARATION INDEX

Elementary chemical composition of marine organisms.
 III. A. P. Vinogradov. *Trav. lab. biogeochem. Acad. Sci. U.R.S.S.* 6, 273 pp. (1944) (in Russian); cf. C.A. 34, 10911. - This concluding vol. of the monograph reviews the metal-contg. blood pigments, the mineralogic compn. of skeletons of marine organisms, and the chem. elements occurring in invertebrates and fishes. A summary of the chem. compn. by types and classes is given. The regulatory influence of the salt mass of ocean water on the chem. compn. of organisms and the basic changes in this compn. during geologic time are discussed. The organisms are indexed by classes. Bibliography. F. L. Gaur.

Trudy Lab. Biogeochem.
 AN SSSR

ASH-ILA METALLOGICAL LITERATURE CLASSIFICATION

GROUPS: A, B, C, D, E, F, G, H, I, J, K, L, M, N, O, P, Q, R, S, T, U, V, W, X, Y, Z, AA, AB, AC, AD, AE, AF, AG, AH, AI, AJ, AK, AL, AM, AN, AO, AP, AQ, AR, AS, AT, AU, AV, AW, AX, AY, AZ, BA, BB, BC, BD, BE, BF, BG, BH, BI, BJ, BK, BL, BM, BN, BO, BP, BQ, BR, BS, BT, BU, BV, BW, BX, BY, BZ, CA, CB, CC, CD, CE, CF, CG, CH, CI, CJ, CK, CL, CM, CN, CO, CP, CQ, CR, CS, CT, CU, CV, CW, CX, CY, CZ, DA, DB, DC, DD, DE, DF, DG, DH, DI, DJ, DK, DL, DM, DN, DO, DP, DQ, DR, DS, DT, DU, DV, DW, DX, DY, DZ, EA, EB, EC, ED, EE, EF, EG, EH, EI, EJ, EK, EL, EM, EN, EO, EP, EQ, ER, ES, ET, EU, EV, EW, EX, EY, EZ, FA, FB, FC, FD, FE, FF, FG, FH, FI, FJ, FK, FL, FM, FN, FO, FP, FQ, FR, FS, FT, FU, FV, FW, FX, FY, FZ, GA, GB, GC, GD, GE, GF, GG, GH, GI, GJ, GK, GL, GM, GN, GO, GP, GQ, GR, GS, GT, GU, GV, GW, GX, GY, GZ, HA, HB, HC, HD, HE, HF, HG, HH, HI, HJ, HK, HL, HM, HN, HO, HP, HQ, HR, HS, HT, HU, HV, HW, HX, HY, HZ, IA, IB, IC, ID, IE, IF, IG, IH, II, IJ, IK, IL, IM, IN, IO, IP, IQ, IR, IS, IT, IU, IV, IW, IX, IY, IZ, JA, JB, JC, JD, JE, JF, JG, JH, JI, JJ, JK, JL, JM, JN, JO, JP, JQ, JR, JS, JT, JU, JV, JW, JX, JY, JZ, KA, KB, KC, KD, KE, KF, KG, KH, KI, KJ, KK, KL, KM, KN, KO, KP, KQ, KR, KS, KT, KU, KV, KW, KX, KY, KZ, LA, LB, LC, LD, LE, LF, LG, LH, LI, LJ, LK, LL, LM, LN, LO, LP, LQ, LR, LS, LT, LU, LV, LW, LX, LY, LZ, MA, MB, MC, MD, ME, MF, MG, MH, MI, MJ, MK, ML, MM, MN, MO, MP, MQ, MR, MS, MT, MU, MV, MW, MX, MY, MZ, NA, NB, NC, ND, NE, NF, NG, NH, NI, NJ, NK, NL, NM, NN, NO, NP, NQ, NR, NS, NT, NU, NV, NW, NX, NY, NZ, OA, OB, OC, OD, OE, OF, OG, OH, OI, OJ, OK, OL, OM, ON, OO, OP, OQ, OR, OS, OT, OU, OV, OW, OX, OY, OZ, PA, PB, PC, PD, PE, PF, PG, PH, PI, PJ, PK, PL, PM, PN, PO, PP, PQ, PR, PS, PT, PU, PV, PW, PX, PY, PZ, QA, QB, QC, QD, QE, QF, QG, QH, QI, QJ, QK, QL, QM, QN, QO, QP, QQ, QR, QS, QT, QU, QV, QW, QX, QY, QZ, RA, RB, RC, RD, RE, RF, RG, RH, RI, RJ, RK, RL, RM, RN, RO, RP, RQ, RR, RS, RT, RU, RV, RW, RX, RY, RZ, SA, SB, SC, SD, SE, SF, SG, SH, SI, SJ, SK, SL, SM, SN, SO, SP, SQ, SR, SS, ST, SU, SV, SW, SX, SY, SZ, TA, TB, TC, TD, TE, TF, TG, TH, TI, TJ, TK, TL, TM, TN, TO, TP, TQ, TR, TS, TT, TU, TV, TW, TX, TY, TZ, UA, UB, UC, UD, UE, UF, UG, UH, UI, UJ, UK, UL, UM, UN, UO, UP, UQ, UR, US, UT, UU, UV, UW, UX, UY, UZ, VA, VB, VC, VD, VE, VF, VG, VH, VI, VJ, VK, VL, VM, VN, VO, VP, VQ, VR, VS, VT, VU, VV, VW, VX, VY, VZ, WA, WB, WC, WD, WE, WF, WG, WH, WI, WJ, WK, WL, WM, WN, WO, WP, WQ, WR, WS, WT, WU, WV, WW, WX, WY, WZ, XA, XB, XC, XD, XE, XF, XG, XH, XI, XJ, XK, XL, XM, XN, XO, XP, XQ, XR, XS, XT, XU, XV, XW, XX, XY, XZ, YA, YB, YC, YD, YE, YF, YG, YH, YI, YJ, YK, YL, YM, YN, YO, YP, YQ, YR, YS, YT, YU, YV, YW, YX, YY, YZ, ZA, ZB, ZC, ZD, ZE, ZF, ZG, ZH, ZI, ZJ, ZK, ZL, ZM, ZN, ZO, ZP, ZQ, ZR, ZS, ZT, ZU, ZV, ZW, ZX, ZY, ZZ.

Ex. 11

1. Unsubstantiated

Chlorine-bromine coefficient of underground waters. A. P. Vinogradov (*Comp. rend. Acad. Sci. U.R.S.S.* 1944, 44, 87-89).
Various factors which bring about variations in the Cl/Br ratio (ρ) of underground waters from the val. (~200) obtained in the analysis of dissolved factors are discussed. Underground waters are subdivided into 3 groups according to the ρ of the water: (1) normal sea H₂O with ρ ~ 200, (2) residual brine of sea salt lakes with ρ < 200, and (3) a solution of NaCl deposits with ρ > 200. C. K. H.

2nd. AN SSSA.

W. I. Vernadsky Lab. Geochem. Problems; Col. Mus. Acad. Sci., 1944.

VINOGRADOV, A. P.

PA 13110

Apr 1946

USSR/Medicine - Goiter
Medicine - Iodine

"Geochemical Conditions in the Regions of Endemic
Goiter," A. P. Vinogradov, 14 pp

"Izv. Ak. Nauk Geograf i Geofiz.", Vol X, No 4


Study of the geochemistry of iodine, and an explanation of the spreading of simple endemic goiter. Data on goiter in the Soviet Union and its connection with the composition of the soil.

13T10

137 AND 7th CODES															100 AND 6th CODES														
PROCEDURES AND PROPERTIES INDEX																													
<div style="font-size: 2em; font-weight: bold; margin-bottom: 10px;">CA</div> <div style="text-align: right; margin-bottom: 10px;">1</div> <div style="text-align: center;"> <p>Vladimir Ivanovich Vernadskii, founder of geochemistry (on the anniversary of his death). A. P. Vinogradov. J. Applied Chem. (U.S.S.R.) 10, 100-112 (1946). A biographical sketch of V.'s life and his achievements in the fields of geochemistry and biogeochemistry, with portrait. M. Schwartz</p> </div> <div style="text-align: right; margin-top: 10px;">Zhur. Prikl. Khim.,</div>																													
<div style="display: flex; justify-content: space-between;"> ASB-154 METALLURGICAL LITERATURE CLASSIFICATION 007742-1429 </div>																													
<div style="display: flex; justify-content: space-between;"> 00000 00000 00000 00000 </div>															<div style="display: flex; justify-content: space-between;"> 00000 00000 00000 00000 </div>														

1ST AND 2ND ORDER		PROCESS AND PROPERTIES INDEX	
CA	COMMON ELEMENTS	<p>A chemical study of the biosphere. A. P. Vinogradov. <i>Podol'sk</i> (U.S.S.R.) 1945, 345-54 (English summary).— V. Vernadskii's concept of the biosphere, the domain of the earth's crust contg. life. Metamorphosed sedimentary rocks down to the crust. foundation are included in the biosphere. In a geochem. study of the biosphere analyses were made of the soils of the Russian plain from north to south along the 40th meridian. The min. and max. contents of rare and other scattered elements in the soil are given in percentage of dry soil, as follows: Li, $1.0-9.1 \times 10^{-4}$; Rb, $0.12-1.7 \times 10^{-4}$; P, $1.0-3 \times 10^{-4}$; Ti, $0.25-0.81$; V, $0.75-2.5 \times 10^{-4}$; Cr, about 5×10^{-4}; Mn, $0.01-0.21$; Cu, $0.12-1.3 \times 10^{-4}$; Ni, $1.0-6.5 \times 10^{-4}$; Co, $0.5-2.6 \times 10^{-4}$; Zn, about 5×10^{-4}; Ga, about 1.0×10^{-4}; Ge, 10^{-4}; As, $1.2-9 \times 10^{-4}$; Se, below 10^{-4}; Br, $0.1-4 \times 10^{-4}$; Rb, $1.0-8.9 \times 10^{-4}$; Sr, $0.013-0.36$; Zr, about 0.03; Mo, $1.5-4 \times 10^{-4}$; Cd, about 2×10^{-4}; I, $0.31-1.7 \times 10^{-4}$; Ba, $0.01-0.30$; Tb, $1.4-2 \times 10^{-4}$; Pb, $0.38-4.3 \times 10^{-4}$; Ra, $1.9-2.9 \times 10^{-4}$; Th, $2.6-9.6 \times 10^{-4}$. Living organisms, as a rule, take up those elements which go into soil. The insol. elements are taken up with difficulty. Thus Ti in soil runs to 0.8%, whereas living organisms contain about 10^{-4} to $10^{-5}\%$; Zr in soil runs up to 0.01%, whereas in organisms it is less than $10^{-4}\%$. A diagram is given showing the elements found in greater quantities in plants and those found in greater quantities in soils. V. theorizes on the relation of the biosphere compn. to the compn. of the living organisms and on the basis of deficiencies and excesses of certain elements he proposes a series of biogeochem. provinces. J. S. Joffe</p>	
		<p>ASB-55A METALLURGICAL LITERATURE</p>	

1st AND 2nd INDEX		3rd AND 4th INDEX	
PROCESSING AND PROPERTIES INDEX			
<p><i>KS</i></p> <p>In memory of Alexander Lvovovich Forman (1903-1945). A. P. Vinogradov. <i>J. Applied Chem. (U.S.S.R.)</i> 19, 608-12 (1946); <i>Chem. Zvest.</i> 1947, 1, 481. M. G. M.</p> <p><i>2</i></p>			
<p>ASSOCIATE METALLURGICAL LITERATURE CLASSIFICATION</p>			
1st AND 2nd INDEX		3rd AND 4th INDEX	
1st AND 2nd INDEX		3rd AND 4th INDEX	



3

Geoschemical history of oxygen and photosynthesis.
A. P. Vinogradov. *Bull. acad. sci. U.R.S.S., Ser. biol.*
1947, 409-22. — A review of various hypotheses on the
geochemistry of O is presented, including the discussion of
the information on O isotope distribution. The isotope O
distribution in photosynthetic O is identical with that of
natural H₂O and differs from that of atm. CO₂, supporting
the idea that photosynthetic O is derived from dehydro-
genation of H₂O and not from CO₂ reduction. The reasons
for the relative abundance of heavy O isotope in the atm.
may be a selective oxidation process series of elements, or
CO₂ in the upper atm. (3. M. Kholodoff

1 — Ak. Nauk SSSR, Ser. Biol.

VINOGRADOV, A. P.

"Boron in the Soils of the Soviet Union,"

Pedology, No. 2, 1947.

Pochvoved.,

PA 1T93

VINOGRADOV, A. P.

USSR/Geochemistry
Minerals - Chemical Properties

1 May 1947

"Isotopic Composition of Oxygen in Certain Minerals,"
A P Vinogradov, E I Dontsova, 2 pp

"Dok Akad Nauk USSR Nov Ser" Vol LVI, No 4

Minerals investigated are cuprite, magnetite, and
chromite.

1T93

CA

112

Determination of the isotopic composition of oxygen due to photosynthesis. A. P. Vinogradov and R. V. Ios. *Compt. rend. acad. sci. U.R.S.S.* 50, 50 (1954) (in English).—On repeating former expts. (C.I. 37, 1954) on the isotopic compn. of O due to photosynthesis in *Elodea canadensis*, V. and T. found that the increase, Δd , in the d. of H_2O prepd. from O obtained by photosynthesis over that of H_2O prepd. from O obtained by electrolysis of water varied from +2.0 to +2.37 (av. +2.2) and differed greatly from the Δd both of O from air ($\Delta d = +7\%$) and from CO_2 ($\Delta d = +11.5\%$). Thus, during photosynthesis O is liberated as a result of H_2O dehydrogenation rather than as a result of the reduction of CO_2 .
Frank Gonet

DOI-AN JSSR,

ASM-SLA METALLURGICAL LITERATURE CLASSIFICATION

CA

8

Isotopic composition of oxygen in aluminosilicates of mineral deposits. A. P. Vinogradov and E. I. Dantsova (V. Vernadskii Geochim. Inst., Leningrad). *Doklady Akad. Nauk S.S.R.* 20, 83-4 (1967). — Examn. of the isotopic O compn. in minerals by heating finally to 2000° with C and bringing the resulting CO into contact with H₂ over Ni-Co catalyst to form H₂O which was analyzed densitometrically showed that the ancient basic and acidic rocks (granite, basalt, diorite, etc.) correspond in their O isotope distribution to that found in water. Hence, no isotope separation apparently took place during the formation of the hard silicate crust and the aq. vapor or liquid phase in the early geol. eras. The apparent isotopic differences that have been found occur only in the biosphere where the exchange processes are proceeding under the attack of atm. O, circulating H₂O, etc. (G. M. Kosolapoff)

CA

Distribution of chemical elements in subterranean waters of various origins. (Significance of coefficients of proportionality.) A. P. Vysotskiy. *Izv. Akad. Nauk SSSR, Geol. Probn., ser. P. P. Sverdlovsk*, 1, 25-33 (1946).—The ratios between chem. elements or ions of similar physico-chem. properties and particularly the ratios between individual chem. elements and their rare or widely-disseminated analogs, elements is used to throw light on the unsolved problems of the genesis of various natural waters (fresh, mineral, and sea). The ratios used are: Ca-Mg, Na-Mg, Na-Mg, Na-Li, K-Li, K-Mg, B-Cl, and Cl-Br. Iodine is exceptional so that the ratios Cl-I and Br-I are not of the same significance for this purpose. The 25 waters whose analyses are listed were collected in 1936 from 3 Russian regions especially for the data. of their content of widely-disseminated chem. elements. Sr and Li were detd. by the T. F. Borovik-Shennikova spectrographic method (C.A. 23, 7695; C.A. 27, 310), with an error of 12%; Br and I, by the method of L. A. Solov'ev and K. V. Menchinskii (C.A. 23, 717). Boron was detd. directly by the method of Tsvetkov (C.A. 24, 6200; 27, 1000; and 40, 6223) with an error of 10%. In addn., many published water analyses contain data for detg. these ratios are quoted, including 5 from Clarke (C.A. 10, 1000). V. H. Gottschalk

CA

15

— Arsenic in the soils of the U.S.S.R. A. P. Vinogradov. *Pedology* (U.S.S.R.) 1948, No. 1, 33-8. Data are given on the As content of 10 profiles which cover practically all the zonal types and some subtypes of soils found in Russia. The quantity of As is fairly uniform. It lies within the limits of 1×10^{-4} to $1 \times 10^{-6}\%$. The average for the soils investigated is $3.5 \times 10^{-6}\%$. The average for the soils from the various parts of the world is $5 \times 10^{-6}\%$. For the soils of the U.S.A. the figure is $7.5 \times 10^{-6}\%$. In regions of recent volcanic activity the As content in the soil is higher. The highest quantity of As is found in the chernozem soils, the lowest in the tundra and podzol soils. The concn. of As takes place in the A horizon of chernozem. In other soils, like the gray semi-desert, the B horizon has the higher quantity of As.

I. S. Ioffe

Pachyved.

J. S. Joffe

AS N-3 LA METALLURGICAL LITERATURE CLASSIFICATION

VINOGRADOV, A. P.

PA 3/49T2

USSR/Academy of Sciences
Chemistry - Analysis

Aug '48

"Works of the Commission on Analytical Chemistry,"
edited by A. P. Vinogradov, Corr Mem, Acad Sci
USSR; reviewed by Prof I. P. Alimarin, Jr Chem Sci,
1 1/2 pp

"Zavod Lab" Vol XIV, No 8.

First volume of series devoted to original articles
and surveys of existing knowledge. Chief defects:
(1) Work of Russian analysts not fully treated;
(2) relative merits of various methods not indicated;
(3) haphazard selection of material. Criticizes
eight articles in detail. Publication of Acad Sci
3 Aug 48

USSR/Academy of Sciences (Contd)

Aug 48

USSR, Moscow, 1948, 195 pp.

3/49T2

VINOGRADOV, A. P.

USSR/Soil Studies
Fluorine

1 Mar 1948

"Fluorine in Soils of the USSR," A. P. Vinogradov,
Corr Mem, Acad Sci USSR; V. V. Danilova, Lab Geochem
Problems imeni V. I. Vernadskiy, 2½ pp

"Dokl Akad Nauk SSSR, Nova-Ser" Vol LIX, No 7

Reports distribution of fluorine in soils of the USSR
Basic source of fluorine in the soils: disintegrated
rocks containing average of 0.03% F, their minerals
containing considerable amount of fluorine, such as
apatite, tourmaline, biotite, muscovite and other
minerals. Gives other secondary sources. Gives
table of fluorine containing soils of the USSR.

47T102

Vinogradova, A. USSR, mss. A. P. Vinogradov and K. G. Vinogradova (*C. N. Acad. Sci., U.S.S.R.*, 1960, no. 897-898). The Mo content of 10 fresh samples from the *Chernobyl* (Minsk) area varies from 1.8 to 12 p.p.m. The values for ash and brown forest soils (Kazan, Tula, Leningrad areas) are 1-4%, for chernozem (Voronezh, N. Caucasus) 1.8-2.8, and for grey chestnut, and red soils (C. Asia, Transcaucasia) 2-4 p.p.m. . Mo Mo content tends to rise with increasing depth of sampling. A. Naumov.

ASME-32A METALLURGICAL LITERATURE CLASSIFICATION

CIA-RDP86-00513R001859910007-1"

VINOGRADOV, A. P.

26376 Mnogokovshevyy zskavator dvoynogo deystviya. (Dobycha gliny). Sbornik
rabot po mest. stroit. Materialam (Upr. prom-sti stroit. Materialov i stroit.
Detaley pri mosgorispolkome, nauch-issled i zksperim. Stantsiya, vyp. 2-3, 1949,
s. 23-26.

SO: LETOPIS' NO. 35, 1949

VINOGRADOV, A. P.

24854. VINOGRADOV, A. P. Biogeokhimicheskiye Provintsiy. Trudy Yubileynoy Sessii,
Posvyashch. Stoletiyu So Dnya Rozhdeniya Dokychayeva. M.-L., 1949, S. 59-84.

SO: Letopis' No. 33, 1949

CA

Chromium and vanadium in the soils of the Soviet-Union. A. P. Vinogradov and G. G. Bergman. *Bykhovskaya* *denie (Pedology)* 1969: 760-773. In soils of the tundra zone there is more V than Cr, probably because of the nepheline syenite. There is more V and Cr in the upper than in the lower horizons. There seems to be more V than in the tundra soils than in many soils of the other soil groups. The soil podzolic soils on clay loams contain more Fe and correspondingly more Cr than the soils on boulder clay. There is more Cr than V in these soils. There is a more or less uniform content of V in the profile. There is a definite concn. of Cr and sometimes of V in the B horizon. In the gray forest soils there is a concn. of V and Cr in the B horizon. Chromium soils contain relatively more Cr and V than the other zonal soils. There is a concn. of Cr and V in the upper horizon, with slightly more Cr. The chestnut-brown and gray semidesert soils have more Cr than V. The red loams, in spite of the high Fe content, contain small amounts of Cr. The V content is higher than in any of the soil groups. In general, the Cr content is higher than that of V. The av. of all soils is Cr 1.9×10^{-4} and V 1×10^{-4} . Tabular data are given on the Cr and V contents of the profiles of the zonal soils in Russia as well as the soils in the U. S., France, England, Germany, and Japan. J. S. Joffe

VINOGRADOV, A.P.

The origin of Urovsik endemic. Trudy Biogeokhim. Lab., Akad. Nauk S.S.S.R.
No.9, 5-29 '49. (MLRA 6:5)
(CA 47 no.16:8200 '53)

VINOGRADOV, A. P.

USSR/Science - Soil Science
Chromium, in soils

Oct 49

"Chromium and Vanadium in Soils of the USSR," A. P. Vinogradov, G. G. Bergman, Lab of Geochem Problems
Imeni V. I. Vernadskiy, Acad Sci USSR, 5 pp

"Pochvoved" No 10

Most soils in USSR lowlands (except tundras and red earth) have an average content of 1.9-1.10-2% chromium and 1.10-2% vanadium. Chromium content in soil horizons is practically the same as that of iron. Vanadium is more or less uniform in distribution. Table gives chromium, vanadium, and iron

1511106

USSR/Science - Soil Science (Contd) Oct 49

content of different soil horizons in various parts of USSR and some foreign countries. Characteristics of chromium and vanadium compounds, especially in humus soils, should be studied. Soluble chromium compounds have been observed, particularly in soils on serpentine rocks.

1511106

VINOGRADOV, A.

24000 VINOGRADOV, A. Shchelochi I.I., Na, K, RE(10s) v pochvakh SSSR. (K uchastii
rasseyannykh v pochvakh Khiz. elementov). Problemy sov. pochvozvedeniya,
SB. 15, 1949, S. 23-30. -- Bibliogr: S. 30.

SO: Letopis, No. 32, 1949.

VINOGRADOV, Aleksandr Pavlovich, 1895-

Geochemistry of scattered rare chemical elements in the soil. Moskva, Izd-vo Akademii nauk SSSR, 1950. 227 p. (52-15919)

QD172.R2V5

1. Earths, Rare

VINOGRADOV, A.

24

113

Fundamental laws of the distribution of trace elements among plants and environment. A. P. Vinogradov. *Mikroelementy v Zhizni Rastenii i Zhivotnykh*, Akad. Nauk S.S.S.R., *Trudy Konf. Mikroelement.* 1950, 7-20 (1952).—A general view of data on distribution of trace elements between the living and the nonliving world. A brief summary is presented of the naturally occurring endemic conditions in plants caused by abnormalities in the supply of the trace elements; a few references to similar effects on the animals are made. Physiological role of trace elements in the plants. M. Ya. Shkol'nik. *Ibid.* 39-54.—A review with numerous references, covering mainly the period 1935-1949, dealing with the effects of trace elements on plant physiology; the elements discussed include Mn, Cu, Zn, B, Mo. Effectiveness of boron fertilizers and conditions of their use. M. V. Katalymov (D. N. Pryanishnikov Agrichim. Exptl. Sta., Dolgoprud). *Ibid.* 78-85; cf. C.I. 45, 1711c.—Effects of traces of B on crops are summarized. Addn. of B to the soil raises the crops of clover and alfalfa, especially in limed soils or those supplied with CaCO_3 ; similar effects are found under field conditions with feed beets, flax, mustard plant, and vegetables in general.

ment of the lacteal system parallels B content. Thus, poppy plants which begin to lose their opium content also lose the viability of the lacteal system; when this phenomenon takes place the content of B in the pods and upper leaves declines significantly. Flower falling in Himalayan *Scopolia* can be prevented by treatment of the bushes with solus. contg. B or by introduction of B into the soil. The normal poppy plants contain 30-54.3 mg. B/kg. of dry plant matter depending on the species. Physiological significance of copper for plants and its effect on crops. M. M. Okuntsov (V. V. Kuibyshev State Univ., Tomsk). *Ibid.* 371-80.—Expts. with addn. of CuSO_4 (0.00045-0.0053%) to cultures of oats, wheat, barley, potato, and beans indicate that the various possible methods of introduction of Cu at moderate concns. aid chlorophyll formation, retard age symptoms, raise protein content, and increase the rate of respiration. Physiological role of copper in plants on peat soils. A. S. Okaneuko and L. E. Ostrovskaya. *Ibid.* 400-9.—Cu deficiency lowers the content of polyphenol oxides, which in turn raises the content of the reduced forms of glutathione and ascorbic acid; at the same time the organism acquires an abnormally low general oxidation-

onev

R. P. Vinogradov

213

reduction potential; this leads to high respiration intensity (50% over control) with exhaustion of sugar supplies. The expts. were made on kok-saghyz plants. Effect of copper on growth and development of potato. N. S. Arkhangel'skaya (Ministry of Agr. U.S.S.R., Moscow). *Ibid.* 410-16.—Introduction of Cu via the extra-radicle route is more effective in the production of beneficial effects of this trace element than is its entry through the soil; spraying or wetting of the plant with CuSO_4 (0.001M) gave up to 18% increase in crop yield. About 12% of added Cu enters the tubers. In peaty soils, wetting of the tubers before planting is the best method of Cu entry. Cu enters mainly in the outer layers of the potato tuber, in sites of eye formation, and highest enzymic activity. Cu spraying or wetting methods lead to increased chlorophyll content of the plant and increased respiration intensity. Molybdenum and its biological role. E. G. Vinogradova (V. I. Vernadskii Inst. Geochem. and Anal. Chem., Acad. Sci. U.S.S.R., Moscow). *Ibid.* 515-38; cf. *C.A.* 46, 7712a.—Re-

view with numerous references covering the period to 1950. The role of molybdenum in the life of plants. A. A. Drobkov (V. I. Vernadskii Inst. Geochem. and Anal. Chem., Acad. Sci. U.S.S.R., Moscow). *Ibid.* 539-45; cf. *C.A.* 46, 7712a.—A brief review in which it is pointed out that traces of Mo are necessary for plants, particularly the leguminous plants. Mo aids the formation of nodules and raises the content of N in the plants. Mo aids the growth of clover and protects it during the winter months, along with increased total plant mass and seed yield. The administration of Mo, as NH_4 molybdate, appeared to be satisfactory at about 1 mg./l. level in the applied soils. The effect of molybdenum on the crop and chemical composition of the bean family plants. I. A. Chernavina. *Ibid.* 640-60.—Addn. of NH_4 molybdate as seed treatment or as soil fertilizer results in improved total crop, seed yield, and root size of the bean family plants. Seed treatment or plant spraying produces good results most economically. The best results are obtained in the year following the use of such mineral supplement. Mo content rises in leaves,

A. P. Vinogradov

33

rootlets, and tubers. Ca content also rises. At the same time the plants display better utilization of N and provide a more intense synthesis of protein matter, with increase of protein to nonprotein N ratio. Leaves of plants supplied with Mo accumulate less sugars than do control plants; this soil carbohydrates are less mobile under conditions of Mo deficiency. Mo also improves P utilization with increase of org. P fraction. Experiments on the effect of boron, molybdenum, and other trace elements on seed sprouting. V. P. Kurnetsov. *Ibid.* 561-7. Seed treatment of cotton seeds with H_2BO_3 soln. (200 mg./l.) for 18 hrs. leads to improved cotton crop by 20% with decrease of pod-falls and hastening of ripening. Solns. of H_2SeO_4 (100 mg./l.) reduces ripening period by 7 days and increases cotton crop by 24%. Treatment with NH_4 molybdate solns. (1000 mg./l.) increases the energy of sprouting; excessive soaking in water leads to a sharp decline in sprouting of the seeds but a soaking in the molybdate soln. for 18 hrs. restores the normal viability of the seeds. Mo stimulates the activity of *Azotobacter*, cocci, and bacteria, particularly in the rhizosphere. Mo aids accumulation of nitrate N in the soil but does not affect P. Mo supplements can increase

cotton crops by about 30%. The effect of some chemical elements on the immunological properties of plants. V. P. Nilova and V. P. Rashevskaya. *Ibid.* 591-592. Seed pretreatment with solns. of trace elements and those needed in larger units, can be used as a means for direction of chemical processes in the plants resulting from the seeds, including the immunological properties. Thus $ZnCl_2$, $NaCl$, and KCl lowered catalase activity in wheat grains and in the resulting sprouts; $KMnO_4$ or $CuCO_3$ raised the activity of catalase and peroxidase. As a result the former seeds are more resistant to brown rust than the latter group. Effects of mineral salt treatment of wheat on general biochem. properties of leaves and crop yield are similarly cited. Summer wheat gives best crop yield increases after seed treatment with $KMnO_4$ or $Ca_3(PO_4)_2$; in winter wheat a similar effect was had from $NaCl$, $ZnCl_2$, $FeCl_3$, H_2BO_3 , and was relatively low with $KMnO_4$ and $Ca_3(PO_4)_2$. No specific data are cited on immunological changes. The role of trace elements in improving the resistance of plants to disease. T. D. Strakhov and T. V. Yuroshenko (State Univ., Kharkov). *Ibid.* 603-12. A supply of trace elements (Cu, Fe, Zn, Ni, Mn) improves the resistance of oats to the attack by *Ustilago avenae*. Fe, Na, and Zn improve the growth activity of the plant. $NaCl$ and $FeSO_4$ in low concns. decreased the size of the *U. avenae* infestations where they did occur. The best results are obtained with Ni and Mn supplied with the normal N, P, and K; Mn is most effective, followed by Ni, Fe, and Zn, in order. Similarly affected is the attack of winter rye by *Urocystis occulta*. G. M. Kosolapoff

CA 8

Change of the chemical composition of the carbonate rocks of the Russian platform. A. P. Vinogradov, A. B. Romanov, and V. M. Ratinskii. *Izvest. Akad. Nauk U.S.S.R., Ser. Geol.* 1952, 22-29.—The qual. changes of chem. compn. of the carbonate rocks of the Russian platform are considered. The trends and periodical character of these changes, and the close connection of the evolution of carbonate compn. with general geotect. changes and the history of the tec-

tonic development of the Russian platform are discussed. Tables of representative chem. analyses are given. Maps are provided to show geographic distribution of the samples. Graphs are given which show (1) Ca and Mg contents vs. time, (2) Ca/Mg content vs. time, (3) Sr and CaSO₄ content vs. time, (4) insol. material and Ba content vs. time, (5) the Th and insol. material content vs. time, and (6) the Ca/Sr content vs. time. Gladys S. Macy.

CA

11 D

Preparation of the products of reduction of carbon dioxide, labelled with carbon¹⁴, by the chloroplasts outside a cell. A. P. Vinogradov, E. A. Bolchenko, and V. I. Beranov. *Doklady Akad. Nauk S.S.S.R.* 78, 327-9(1951); cf. *C.A.* 44, 7937h. — Chloroplasts from white clover or primula, isolated from the cells, were examd. in CO₂ labelled with C¹⁴ in the presence of 2% O by manometric technique. Control expts. with boiled chloroplasts gave no C¹⁴ activity in the exts., but active chloroplasts gave significant activity, and almost all activity was pptd. by Ba as a complex, which contains 0.8% P and is free of N, consisting largely of carbohydrate-like materials, giving uronic acid reaction, and other carboxylic acids. The latter increase in proportion on cloudy days, the former predominate in specimens taken on sunny days. In prolonged expts. some 70% of the retained C¹⁴ is extractable with EtOH, in the ketone fractions. Thus, the process of reduction proceeds via carboxylic acids which do not reduce Cu, then uronic acids, then ketones. (I. M. Konolapoff

VINOGRADOV, A.P., ZADOROZHNYI, I.K., ZYKOV, S.I.

Lead - Isotopes

Composition of lead isotopes and the age of the earth. Dokl. AN SSSR 85, No. 5, 1952.

Monthly List of Russian Accessions, Library of Congress, December 1952. Unclassified.

VINCORADOV, A. P.; DONTSOVA, YE. I.

Oxygen - Isotopes

Isotopic oxygen constituent of minerals of scarn origin. Dokl. AN SSSR 85 no. 6, 1952.

9. Monthly List of Russian Accessions, Library of Congress, December 195~~8~~₂, Unclassified.

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859910007-1

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859910007-1"

VINOGRADOV, A. P.

Vinogradov, A. P. Geochemie seltener und nur in Spuren
vorhandener chemischer Elemente in Böden. Berlin: Akad.
Verlag. 1964. 230 pp. DM 27.

Translation of Title: Geochemistry of
Rare and Trace Elements in Soils.

we

ge

VINOGRADOV, A. P.

USER/ Geology - Geochemistry

Card 1/1 Pub. 46 - 1/19

Authors : Vinogradov, A. P.

Title : The geochemistry of isotopes

Periodical : ^{Ak. Nauk} Izv. AN SSSR. Ser. geol. 3, 3 - 19, May - Jun 1954

Abstract : Excerpts are presented of the lecture entitled, "The Geochemistry of Isotopes," given by the author on February 1, 1954 during the general meeting of the Academy of Sciences, USSR. It is pointed out that by studying the isotopic composition of elements in litho-, atmo-, and hydrospheres it would be possible to study the following important geological problems: development of the earth as a whole and individual fragment of the earth's crust, origin of the atmosphere, problems of diffusion of oxygen, sulfur, carbon, hydrogen in the crust of the earth and the connection between the hypogeneus and hypergeneus processes occurring in the earth, and so on. Tables; graphs; drawings; illustration.

Institution:

Submitted: February 5, 1954

VINOGRADOV, A. P.

USSR.

✓ Search for ore deposits by means of plants and soils.
A. P. Vinogradov. *Trudy Biogekhim. Lab., Akad. Nauk
S.S.S.R.* 18, 3-27 (1954); *ibid.* 19, 438g. — A detailed
review, with numerous references, of the methods used to
locate ores by chem. analysis for the desired metals of the
plants and soils in the area; plants in the ore location usually
show metal enrichment by a factor of 10 or greater over nor-
mal levels. G. M. Kosolapoff

ex ju

VINOGRADOV, A. P.

USSR/Miscellaneous - Isotopes

Card : 1/1

Authors : Vinogradov, A. P., Academician

Title : Geochemistry of isotopes

Periodical : Vest. AN SSSR, 24, Ed. 5, 26 - 43, May, 1954

Abstract : A general description of various methods for the determination of terrestrial and atmospheric ages, made possible by determining the changes in isotopic ratios of the elements. Calls special attention to a method based on the determination of the C^{14} isotope in the atmosphere, and illustrates an installation for detecting the C^{14} in the atmosphere. Charts and tables showing variations with the ages of such isotopic ratios as O^{16}/O^{18} , C^{12}/C^{13} and S^{32}/S^{34} . The possibility of determining the role which these isotopes played in formation of rocks and minerals is also discussed.

Institution : ...

Submitted : ...

VINOGRADOV, A.P.

[Physicochemical control methods of uranium production] Fiziko-khimicheskie metody kontrolya proizvodstva urana; doklady, predstavlenyye SSSR na Mezhdunarodnuyu konferentsiyu po mirnomu ispol'zovaniyu atomnoy energii. Moskva, 1955. 20 p. [Microfilm]
(Uranium) (MIRA 9:3)

VINOGRADOV, A.P., akademik, redaktor.

[Use of tagged atoms in analytical chemistry] *Primenenie mechenykh atomov v analiticheskoy khimii*. Moskva, Izd-vo Akademii nauk SSSR, 1955. 233 p. (MLRA 8:11)

1. Akademiya nauk SSSR. Institut geokhimii i analiticheskoy khimii.
(Chemistry, Analytical) (Radioactive tracers)

VII. Sbornik 51.
STARIK. I.Ye., redaktor; SHCHERBAKOV, D.I., akademik, redaktor; VINOGRADOV, A.P., akademik, redaktor; POLKANOV, A.A., akademik, redaktor; SHATSKIY, N.S., akademik, redaktor; BARANOV, V.I., professor, redaktor; PEKARSKAYA, T.B., kandidat geologo-mineralogicheskikh nauk, redaktor; CHERDYNTSEV, V.V., redaktor; POLYAKOVA, T.V., tekhnicheskij redaktor.

[Transactions of the third session of the Committee for Determining the Absolute Age of Geological Formations, March 25-27, 1954] Trudy tret'ei sessii, 25-27 marta 1954. g. Moskva, 1955. 260 p. [Microfilm] (MLRA 9:1)

1. Akademiya nauk SSSR. Komissiya po opredelniya absolyutnogo vozrasta geologicheskikh formatsii. 2. Chlen-korrespondent AN SSSR (for Starik). (Geological time)

BARANOV, V.I.; VINOGRADOV, A.P., akademik, redaktor; MYASNIKOV, I.A.
redaktor; STRUCHKOV, Yu.T., redaktor; MOSKVICHEVA, N.I., tekhnicheskiiy redaktor.

[Radiometry] Radiometriia. Moskva, Izd-vo Akademii nauk SSSR,
1955. 327 p. (MLRA 8:12)
(Radiation--Measurement)

91.

A D

... has been observed the

"APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859910007-1

APPROVED FOR RELEASE: 09/01/2001

CIA-RDP86-00513R001859910007-1"

VERNADSKIY, Wladimir Ivanovich, 1863-1945; VINOGRADOV, A.P., akademik,
redaktor; SHUBNIKOVA, O.M., doktor geologo-mineral'nykh nauk;
redaktor; KUN, H.P., redaktor; ZELENIKOVA, Ye.V., tekhnicheskiy
redaktor.

[Selected works] Izbrannye sochineniia. Moskva, Izd-vo Akademii
nauk SSSR, Vol.2, 1955. 615 p. (MLRA 8:12)
(Geochemistry) (Mineralogy)

FERSMAN, A.Ye., akademik; BELOV, N.V., akademik, redaktor; VINOGRADOV,
A.P., akademik, redaktor; SHCHERBAKOV, D.I., akademik, redaktor;
KAPUSTINSKIY, A.P., redaktor; SAUKOV, A.A., redaktor; SHCHERBINA,
V.V., doktor geologo-mineral'nykh nauk, redaktor; KUM, N.R.,
redaktor; MOSKVICHEVA, N.I., tekhnicheskiy redaktor.

[Selected works] Izbrannye trudy. Moskva, Izd-vo Akademii nauk SSSR.
Vol.3. 1955. 798 p. (MLRA 8:12)

1. Chlen-korrespondent AN SSSR (for Kapustinskiy and Saukov)
(Geochemistry)

VINOGRADOV, A. P.

✓ Radiochemical studies of the products of nuclear transformations under bombardment with high-energy particles. *NU*
- A. P. Vinogradov. *Sessiya Akad. Nauk SSSR po Mirovym Atomnoi Energii* 1953, Plenarное Zasedanie, 60-78 (English summary, 78-80).—A summary of work done during 1950-2 on nuclear transformations in U, Th, Bi, Ag, Cu, W, and La caused by high-energy protons, neutrons, and He^{++} at 230-680 m.e.v. Radioisotope distribution curves are presented and the studies extended to medium and light nuclei. 13 references. G. M. K.

VINOGRADOV, A.P.

USSR/Optics - Optical Methods of Analysis. Instruments.

K-7

Abs Jour : Referat Zhur - Fizika, No 3, 1957, 7948

Author : Vinogradov, A.P.

Title : Physico-Chemical Methods of Control of Production of Uranium.

Orig Pub : I ssledovaniya v obl. geologii, khimii. i metallurgii. M.,
Izd-vo AN SSSR, 1955, 72-89

Abstract : Survey of methods for determining U as applied to its control of manufacture. A brief description is given for the spectral methods of quantitative determination of impurities in U; the determination of small quantities of U in rocks, ores, and semi-finished articles; method of isotopic addition in the spectral determination of U in ores and semi-finished products; X-ray spectral method of determining U and of its impurities; and polarographic determination of U.

Card 1/1

- 97 -

Vinogradov, A.P.

6
12ML

4039 AEC-47-2435(PL 2) (p. 187-204)
LEAD ISOTOPES AND THEIR GEOLOGICAL SIGNIFICANCE.
A. P. Vinogradov, p. 187-204 of CONFERENCE OF THE
ACADEMY OF SCIENCES OF THE USSR ON THE PEACE-
FUL USE OF ATOMIC ENERGY, JULY 1-6, 1955.
SESSION OF THE DIVISION OF CHEMICAL SCIENCE.

(Translation). 20p.

This paper was originally abstracted from the Russian
and appeared in Nuclear Science Abstracts as NSA 9-7776.

RSJ PMZ JPH

VINOGRADOV, A.P.

The development of geochemistry. Vest. Mosk. un. lp no. 45:
169-184 Ap-May '55. (MIRA 8:8)
(Geochemistry)